CLAIMS

1. An immunoassay for detecting exposure to *Leishmania* parasites in a subject comprising the steps of:

contacting a sample from the subject suspected of having leishmaniasis with a soluble antigen prepared by utilizing a protein-free medium; and

detecting the presence or measuring the amount of an antibody or fragment thereof in the sample bound to the soluble antigen.

- 2. The immunoassat of claim 1 wherein the protein-free medium comprises D, xylose.
- 3. The immunoussay of claim 1 wherein the protein-free medium further comprises at least one of the following ingredients: Hepes buffer, L-glutamine and sodium bicarbonate without phonol red.
- 4. The immunoassay of claim 1 wherein the antibody is IgG or IgM and is specific for a Leishmania antigen.
 - 5. The immunoassay of claim wherein the sample is a serum sample.
- 6. The immunoassay of claim 5 wherein the serum sample is modified by diluting it 1:1000 in blocking buffer having 1.0% boiled casein.
- 7. The immunoassay of claim 1 wherein said immunoassay is capable of diagnosing visceral, cutaneous or canine leishmaniasis in a subject.
- 8. The immunoassay of claim 1 wherein the *Leishmania* soluble antigen preparation is prepared by using clones of *Leishmania donovani* or *Leishmania mexicana*.
- 9. An immunoassay for diagnosing leishmaniasis in a subject comprising the steps of:

5

20

115500

25

30





contacting a sample from the subject with an antibody or fragment thereof that specifically binds the *Leishmania* exo-antigen; and

detecting the presence or measuring the amount of said antibody or fragment thereof bound to said Leishmania exo-antigen.

5

10. The immunoassay of claim 9 wherein the antibody or fragment thereof is adsorbed onto a substrate.

10

11. A kit for the diagnosis of leishmaniasis in a subject comprising a substrate and a soluble antigen of either L. donovani or L. mexicana prepared by utilizing a protein-free medium packaged together for multiple or single use assays.

12. The kit of claim 1 wherein the substrate is coated with the soluble antigen.

The kit of claim 11 further comprising a positive control. 13.

14. The kit of claim 11 further comprising a negative control.

15.

16.

19.

20.

to a label.

The kit of claim 11 further comprising a diluent.

25

17. The kit of claim 11 further comprising a substrate chromogen.

The kit of claim 11 further comprising an anti-human IgG conjugated

- 18.
- The kit of claim 11 further comprising a substrate buffer.

The kit of claim 11 further comprising a blocking buffer.

The kit of claim 11 further comprising a stopping solution.

30

5

- 21. A kit for the detection of an exo-antigen of a leishmania parasite in a sample comprising a fluorescein-labeled antibody that binds the exo-antigen and a counter-tain packaged together for multiple or single use assays.
- 22. The kit of claim 21 further comprising a protein stabilized buffer solution.
 - 23. The kit of claim 21 further comprising sodium azide.
- 24. The kit of claim 22 wherein the fluorescein-labeled antibody is diluted in the protein stabilized buffer saturion.
 - 25. The kit of claim 21 wherein the counter-stain is Evans Blue.
- 26. The kit of claim 21 further comprising a solid substrate to which the sample to be tested is fixed.
 - 27. The kit of claim 1 further comprising instructions.
 - 28. The kit of Jaim 21 further comprising instructions.
- 29. A diagnostic device comprising a *Leishmania* soluble antigen prepared by utilizing a protein-free medium and a means for detecting an antibody bound to the *Leishmania* soluble antigen.
- 30. A diagnostic device comprising an antibody or fragment thereof that binds an exo-antigen found in a conditioned medium made by cultivating a *Leishmania* parasite in a protein-free medium.
- 31. A method of preparing a diagnostic device comprising adsorbing to a substrate an antibody or fragment thereof which binds an exo-antigen found in conditioned medium made by cultivating a *Leishmania* parasite in a protein-free medium.

5

H

- 32. A method of preparing a diagnostic device comprising adsorbing to a substrate a soluble antigen of a *Leishmania* parasite prepared by utilizing a proteinfree medium.
- 33. A method of detecting a *Leishmania* parasite in a sample comprising contacting the sample with an antibody specific for an exo-antigen found in a conditioned medium made by cultivating the *Leishmania* parasite in a protein-free medium; and

detecting the presence or measuring the amount of an antibody or fragment thereof in the sample bound to an antigen in the sample.

- 34. A protein-free medium comprising an agent that balances the oncotic pressure across a semi-permeable cell membrane.
 - 35. The protein-free medium of claim 34, wherein the agent is D, xylose.
- 36. The protein-free medium of claim 34, further comprising at least one of the following ingredients: Hepes buffer, L-glutamine and sodium bicarbonate without phenol red.
- 37. A method for obtaining an exo-antigen from an organism comprising culturing the organism in a protein-free medium.
 - 38. The method of claim 37, wherein the organism is a leishmania parasite.
 - 39. The method of claim 37, wherein the protein-free medium is XOM.
 - 40. The exo-antigen obtained according to the method of claim §7.